




SIGNIFICANTLY **REDUCE** RECTAL **SIDE EFFECTS** of prostate radiation therapy¹

with Barrigel™ Rectal Spacer



WORRIED ABOUT **RECTAL SIDE EFFECTS** FROM PROSTATE RADIATION THERAPY?

Being located so close to the prostate puts the rectum at risk for high-dose radiation exposure during treatment.

This can result in complications such as (but not limited to):²



PAIN



RECTAL SWELLING



HEMORRHOIDS



CHRONIC DIARRHEA

Barrigel™ rectal spacer helps protect the rectum by reducing the amount of high-dose radiation that reaches it during treatment.¹

It is a body-safe gel that is placed between the prostate and rectum before radiation therapy starts.^{3,4}

CHOSEN BY PATIENTS
& THEIR DOCTORS
FOR OVER

75,000^{††}


SPACING
PROCEDURES

**PROVEN SAFE
& EFFECTIVE**

Barrigel rectal spacer:

- Made from hyaluronic acid, which is found naturally in the body^{5,6,†}
- Biocompatible and naturally absorbed by the human body^{1,3,5}
- Creates safe, consistent space between the prostate and rectum^{1,5,7,8}
- Clinically proven to reduce rectal side effects of prostate radiation therapy¹

THE IMPORTANCE OF SPACE



Barrigel spacer creates space between the prostate and rectum, moving rectal tissue away from the high-dose radiation area.¹

The spacer maintains its shape and separation for the duration of your radiation therapy.^{1,5} Your body absorbs the biocompatible gel over time.⁵

Visit Barrigel.com/procedure to learn more about the spacing procedure.

WHY DOCTORS CHOOSE BARRIGEL RECTAL SPACER

- Clinically proven to significantly reduce unwanted rectal radiation exposure in 98.5% of patients¹
- Proven superior in the reduction of acute grade 2+ GI toxicity^{1,3,9,10}
- The only rectal spacer that can be sculpted to fit the unique contours of your anatomy^{1,3,5,7,8}
- Zero reports of discomfort or rectal fullness post-procedure in their pivotal study^{1,3,9,10}



After doing the radiation with Barrigel [spacer] I had no rectal side effects."

Michael B.

Prostate Cancer Survivor
Individual results may vary



To learn more, scan this code or visit Barrigel.com/Resources



Barrigel™

References

1. Mariados NF, Orio PF III, King MT et al. *JAMA Oncol* (2023).*
2. American Cancer Society. Radiation Therapy for Prostate Cancer. Accessed April 2025. <https://www.cancer.org/cancer/prostate-cancer/treating/radiation-therapy.html>
3. Data on File. As of 10/01/2024.
4. Barrigel Injectable Gel Instructions for Use (2022).
5. Svatos M, Chell E, Low DA et al. *Med Phys* (2024)*
6. Restylane® celebrates 25 years of natural-looking results with its signature line of hyaluronic acid fillers. 2021
7. Williams J, Mc Millan K, Chao M et al. *J Med Imag Radiat Sci* (2022).
8. Gejerman G, Goldstein MM, Chao M et al. *Pract Radiat Oncol* (2023).**
9. Mariados N, Sylvester J, Shah D et al. *Int J Radiat Oncol Biol Phys* (2015).
10. Song D et al. *Int J Radiat Oncol Biol Phys* (2024).

*Study sponsored by Palette Life Sciences, now part of Teleflex.

**Drs Gejerman, Chao, Lederer, and Orio are paid consultants of Palette Life Sciences, now part of Teleflex

†Dr. Klaassen is a paid consultant of Teleflex

¹Barrigel, made from NASHA, is indicated for rectal spacing in prostate cancer patients and was FDA-cleared in May 2022.

¹Management estimate based on product sales as of: October 2025

Barrigel™ rectal spacer is intended to temporarily position the anterior rectal wall away from the prostate during radiotherapy for prostate cancer and, in creating this space, it is the intent of Barrigel rectal spacer to reduce the radiation dose delivered to the anterior rectum. Barrigel rectal spacer is composed of biodegradable material and maintains space for the entire course of prostate radiotherapy treatment and is intended to be absorbed by the patient's body over time. Barrigel rectal spacer should only be administered by qualified and properly trained physicians with experience in ultrasound guidance and injection techniques in the urogenital/pelvic area. As with any medical treatment, there are some risks involved with the use of Barrigel rectal spacer. Potential complications associated with the use of Barrigel rectal spacer include, but are not limited to: pain associated with Barrigel rectal spacer injection; needle penetration of the bladder, prostate, rectal wall, rectum, or urethra; injection of Barrigel rectal spacer into the bladder, prostate, rectal wall, rectum, urethra, or intravascularly; local inflammatory reactions; infection; urinary retention; rectal mucosal damage, ulcers, necrosis; bleeding; constipation; and rectal urgency. Discuss these adverse effects with your healthcare provider. For more information about Barrigel rectal spacer, please visit [barrigel.com](https://Barrigel.com). Individual results may vary.

Caution: Federal (USA) law restricts this device to sale by or on the order of a physician.

Teleflex, and the Teleflex logo, Barrigel, and the Barrigel logo are trademarks or registered trademarks of Teleflex Incorporated or its affiliates, in the U.S. and/or other countries.

© 2026 Teleflex Incorporated. All rights reserved. APM1222A



Barrigel spacer is the only rectal spacer I use in my practice...based on my experience, the data, patient safety, and results we've seen."

Zach Klaassen

MD, MSc[‡]

Urologic Oncologist

